

# Does Lyndon's Lenth Function Imply Imply the Universal Theory of Free Groups?

ANTHONY M. GAGLIONE<sup>1</sup> and DENNIS SPELLMAN

ABSTRACT. This note shows that every model of the universal theory of the non-Abelian free groups admits a Lyndon length function. The question is then posed as to whether the model class of the universal theory of the non-Abelian free groups is precisely the class of such groups. The authors have subsequently given a negative answer to this question.

Definitions and notation will be that of Bell and Slomson [3], and of Gaglione and Spellman [6], [7] and [8].

Let  $L$  be a first-order language with equality. Two  $L$ -structures  $A$  and  $B$  *have the same universal theory* just in case they satisfy precisely the same universal sentences (and therefore also precisely the same existential sentences) of  $L$ . If  $B$  is an  $L$ -structure, let  $\text{Th}(B) \cap (\forall \cup \exists)$  be the set of all universal and existential sentences of  $L$  true in  $B$ . Evidently the  $L$ -structure  $A$  has the same universal theory as  $B$  if and only if  $A$  is a model of  $\text{Th}(B) \cap (\forall \cup \exists)$ .

If  $A$  is a substructure of the  $L$ -structure  $B$ , then a necessary and sufficient condition that  $A$  and  $B$  have the same universal theory is that there be a model  $*A$  of

$\text{Th}(A) \cap (\forall \cup \exists)$  such that  $A \subseteq B \subseteq *A$ . This in turn is equivalent to the existence of an index set  $I$  and an ultrafilter  $D$  on  $I$  such that  $B$  is embeddable in the ultrapower  $A^I/D$ . A different necessary and sufficient condition that an  $L$ -structure  $B$  and a substructure  $A$  have the same universal theory is that  $A$  and  $B$  satisfy precisely the same primitive sentences of  $L$ . (See [3], Ch.9.)

Let  $A$  be a non-trivial, torsion-free, Abelian group. Let  $<$  be a strict linear order on  $A$  such that for arbitrary  $(a, b, c) \in A^3$  we have  $a + c < b + c$  whenever  $a < b$ . Then the ordered pair  $\Lambda = (A, <)$  is an *ordered Abelian group*. A non-trivial, torsion-free, Abelian group  $A$  is *orderable* provided there is at least one strict linear order  $<$  such that  $\Lambda = (A, <)$  is an ordered Abelian group. It is well-known that every non-trivial, torsion-free, Abelian group is

---

<sup>1</sup>1991 Mathematics Subject Classification. Primary 20A15, 20F32, 03C60. Secondary 03B10, 03C07, 03C20, 20E08. ■

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\TeX}$

<sup>1</sup>The research of this author was partially supported by the Identification Systems Branch, Radar Division, Naval Research Laboratory.

orderable. None the less, we'll present an argument to that effect in this paper. We shall also show the well-known result that this class of Abelian groups is the class having the same universal theory as  $\mathbb{Z}$ ; moreover, we show that every model of the universal theory of the non-Abelian free groups admits a Lyndon length function.

We now specify two first-order languages with equality.  $L_o$  shall contain a binary operation symbol  $\cdot$ , a unary operation symbol  $^{-1}$  and a constant symbol  $1$ .  $L$  shall contain a binary operation symbol  $+$ , a unary operation symbol  $-$ , a constant symbol  $0$  and a binary relation symbol  $<$ .  $L_o$  shall be the *language of group theory* and  $L$  shall be the *language of ordered Abelian groups*. A primitive sentence of  $L_o$  is one of the form

$$\exists(\mathbb{I}(p_i()) = P_i())\mathbb{I}(q_j() \neq Q_j()))$$

where  $\bar{x}$  is a tuple of variables and the  $p_i(x), P_i(x), q_j(x)$  and  $Q_j(x)$  are terms of  $L_o$ .

In case the  $L_o$ -structures we are considering are groups this type of sentence may be simplified to one of the form

$$\exists(\mathbb{I}(p_i() = 1)\mathbb{I}(q_j() \neq 1))$$

where  $\bar{x} = (x_1, \dots, x_m)$  is a tuple of distinct variables and the  $p_i(\bar{x})$  and  $q_j(\bar{x})$  are words on  $\{x_1, \dots, x_m\} \cup \{x_1^{-1}, \dots, x_m^{-1}\}$ .

LEMMA 1. *A non-trivial Abelian group A has the same universal theory as  $\mathbb{Z}$  if and only if A is torsion-free.*

PROOF. One implication is trivial. Assume A is a non-trivial, torsion-free, Abelian group to show that A is a model of

$\text{Th}(\mathbb{Z}) \cap (\forall \cup \exists)$ . We write our groups additively here. Let  $a$  be a non-zero element of A and put  $A_0 = \langle a \rangle \cong \mathbb{Z}$ . It will suffice to show that  $A_0$  satisfies every primitive sentence true in A. To that end consider the system

$$(*)$$

of equations and inequations. Suppose  $(*)$  has a solution  $(x_1, \dots, x_m) = (a_1, \dots, a_m)$  in A. It suffices to show that  $(*)$  has a solution in  $A_0$ . Let B be

the subgroup of  $A$  generated by  $\{a_1, \dots, a_m\}$ . If  $B = 0$ , then  $(x_1, \dots, x_m) = (0, \dots, 0)$  is a solution to  $(*)$  in  $A_0$ . We may therefore assume  $B \neq 0$ . So  $B$  is then a non-trivial, finitely generated, torsion free, Abelian group. Thus,  $B$  is free Abelian of some finite rank  $r \geq 1$ . But it was shown in Gaglione and Spellman [7] that  $\mathbb{Z}^r$  and  $\mathbb{Z}$  have the same universal theory. Therefore  $(*)$  has a solution in  $A_0$ . ■

DEFINITION 1 (Lyndon [12]). Let  $G$  be a (multiplicatively written) group. Let  $\Lambda = (A, <)$  be an (additively written) ordered Abelian group. Let  $\lambda : G \longrightarrow A, g \mapsto |g|$  be a function and let  $2c : G^2 \longrightarrow A$  be defined by  $(g_1, g_2) \mapsto |g_1| + |g_2| - |g_1 g_2|$ . The ordered triple  $(G, \Lambda, \lambda)$  is a *normed group* provided the following six axioms are satisfied:

- (A0)  $x \neq 1$  implies  $|x| < |x^2|$
- (A1)  $|x| \geq 0$  and  $|x| = 0$  iff  $x = 1$
- (A2)  $|x^{-1}| = |x|$
- (A3)  $2c(x, y) \geq 0$
- (A4)  $2c(x, y) > 2c(x, z)$  implies  $2c(y, z) = 2c(x, z)$
- (CO)  $2c(x, y) \equiv 0 \pmod{2A}$

A group  $G$  is *normable* provided there is at least one ordered Abelian group  $\Lambda = (A, <)$  and at least one map  $\lambda : G \longrightarrow A$  such that  $(G, \Lambda, \lambda)$  is a normed group.

REMARK. The axioms are not independent. Chiswell [5] has shown that (A3) is a consequence of (A2), (A4) and the following (A1')  $|1| = 0$ .

If  $S$  is a set of sentences of a first-order language  $L$  with equality, let  $\mathbb{M}(S)$  be the model class of  $S$ .

THEOREM 1. *Let  $L$  be a first-order language with equality. Let  $X$  be a class of  $L$ -structures. Then the following three properties form a set of necessary and sufficient conditions that  $X$  be of the form  $\mathbb{M}(S)$  for at least one set  $S$  of sentences of  $L$ :*

- (i)  *$X$  is closed under isomorphism.*
- (ii)  *$X$  is closed under the formation of ultraproducts.*

(iii) *If  $cX$  is the class of all  $L$ -structures not in  $X$ , then  $cX$  is closed under the formation of ultrapowers.*

Theorem 1 is a deduction of Theorem 3.10, Chapter 7 of [3] without assuming (G.C.H.) using the Keisler-Shelah Theorem. (See [3] and [15].)

Although "the" norm is not generally defined in  $L_o$ , norms are internal in the sense that they extend to ultraproducts. It is then straightforward to deduce -

**COROLLARY.** *The class of all non-Abelian, normable groups is the model class  $\mathbb{M}(\Theta)$  of some set  $\Theta$  of sentences of  $L_o$ .*

It is known that the non-Abelian free groups have the same universal theory (see [7]). Thus, if  $F_2$  is free of rank 2 and  $\Phi = \text{Th}(F_2 \cap (\forall \cup \exists))$ , then every non-Abelian, free group is a model of  $\Phi$ . It is not difficult to convince oneself that the models of  $\Phi$  are precisely those non-Abelian groups embeddable in some ultrapower of  $F_2$ , since one can easily show that every model of  $\Phi$  contains a copy of  $F_2$ . But if  $F_2 = \langle a_1, a_2 \rangle$ , then the length function with respect to the free basis  $\{a_1, a_2\}$ ,

$\lambda : F_2 \longrightarrow \mathbb{Z}$  induces a length function  $*\lambda : F_2/D \longrightarrow \mathbb{Z}^I/D$  making  $(F_2/D, (\mathbb{Z}^I/D, <), *\lambda)$  into a normed group whenever  $I$  is an index set and  $D$  is an ultrafilter on  $I$ . The restriction of  $*\lambda$  to the subgroup  $G$  makes  $G$  into a normed group. Thus, every model of  $\Phi$  is also a model of  $\Theta$ . In symbols -

**THEOREM 2.**  $\mathbb{M}(\Phi) \subseteq \mathbb{M}(\Theta)$ .

Brignole and Ribeiro [4] have given a proof of a theorem of Gurevic and Kokorin asserting that any two ordered Abelian groups have the same universal theory. Since any ordered Abelian group  $\Lambda = (A, <)$  contains an ordered subgroup isomorphic to  $(\mathbb{Z}, <)$  it follows that every ordered Abelian group  $\Lambda$  is embeddable in some ultrapower

$$(\mathbb{Z}, <)^I/D = (\mathbb{Z}^I/D, <).$$

Thus, every normed group admits a norm with values in an ordered Abelian group of the form  $(\mathbb{Z}^I/D, <)$ .

Let  $L$  be a first-order language with equality. A sentence of  $L$  of the form  $\forall \bar{x} \exists \bar{y} \phi(\bar{x}, \bar{y})$  where  $\bar{x}$  and  $\bar{y}$  are disjoint tuples of variables,  $\phi(\bar{x}, \bar{y})$  contains no quantifiers and  $\phi(\bar{x}, \bar{y})$  contains free at most the variables in  $\bar{x}$  and  $\bar{y}$  is a *universal-existential* sentence of  $L$ . Any sentence of  $L$  logically equivalent to a universal-existential sentence of  $L$  is a  $\mathbf{a}\mathbf{b}_2$ -sentence of  $L$ . Since vacuous quantifications are permitted, every universal sentence of  $L$  and every existential sentence of  $L$  is also a  $\mathbf{a}\mathbf{b}_2$ -sentence of  $L$ .

**THEOREM 3.**  $\Theta$  may be taken to be a set of  $\mathbf{ab}_2$ -sentences of  $L_o$ .

**PROOF.** In view of Theorem 2, p. 279 of Grätzer [10], it suffices to show that the union  $G = \bigcup_{n < \omega} G_n$  of a chain  $(G_n)_{n < \omega}$  of non-Abelian, normable subgroups  $G_0 \subseteq G_1 \subseteq \dots \subseteq G_n \subseteq \dots$  is normable. To that end suppose that  $(G_n, \Lambda_n, \lambda_n)$  is a normed group. Let  $D$  be a non-principal ultrafilter on  $\omega$  and let  $\Lambda =$

$(\prod_{n < \omega} \Lambda_n) / D$  be the ultraproduct of the family  $(\Lambda_n)_{n < \omega}$  of ordered Abelian groups with respect to the ultrafilter  $D$ . Let  $\Lambda = (A, <)$ . For each  $g \in G$ , let  $\deg(g) = \min\{n \in \omega \mid g \in G_n\}$ . Finally, let

$\lambda : G \longrightarrow A$  be given by  $g \mapsto L_g / D$  where

Then it is straightforward to verify that  $(G, \Lambda, \lambda)$  is a normed group. ■

**QUESTION.** Is  $\mathbb{M}(\Phi) = \mathbb{M}(\Theta)$  ?

Equivalently: *Does every non-Abelian, normable group have the same universal theory as the non-Abelian free groups?*

In view of (5.3), (5.4)(6.4) of Alperin and Bass [1], we may also pose the

**QUESTION.** *Let  $G$  be a non-Abelian group. Is it the case that  $G$  is a model of  $\Phi$  if and only if there is an ordered Abelian group  $\Lambda$  and a  $\Lambda$ -tree  $T$  such that  $G$  acts freely on  $\Lambda$  without inversions?*

(i.e., if and only if  $G$  is tree-free in the sense of Bass - see [2].)

#### Addendum

Since the original preparation of the manuscript, the authors have learned of [14]. In that work, Remeslennikov also shows that every model of  $\Phi$  is

normable. Moreover, a negative answer to our question is given independently in [9] and in [14].

The authors wish to thank the referee for helpful comments.

#### REFERENCES

- (1) Alperin, R., and Bass, H., "Length functions of group actions on  $\Lambda$ -trees," *Annals of Mathematical Studies, Combinatorial Group Theory and Topology*, Gersten, S. M. and Stallings, J. R., Editors, Princeton University Press, Princeton, NJ(1987), 265 – 378.
- (2) Bass, H., "Group actions on non-Archimedean trees," *Mathematical Sciences Research Institute Publications, Arboreal Group Theory*, Alperin, R. C., editor, Springer-Verlag, New York, NY(1991), 69 – 131.
- (3) Bell, J. L. and Slomson, A. B., *Models and Ultraproducts*, (Second revised printing), North- Holland, 1971, Amsterdam.
- (4) Brignole, D. and Ribeiro, H., "On the universal equivalence for ordered abelian groups," *Algebra i Logika Sem.* 4 (1965), no. 2, 51 – 55.
- (5) Chiswell, I. M., "Abstract length functions in groups," *Math. Proc. Camb. Phil. Soc.* 80(1976), 451 – 463.
- (6) Gaglione, A. M. and Spellman, "More model theory of free groups," Preprint.
- (7) Gaglione, A. M. and Spellman, D., "Some model theory of free groups and free algebras," *Houston J. Math.* to appear.
- (8) Gaglione, A. M. and Spellman, D., "Even more model theory of free groups," *Proc. of AMS Conference on Infinite Groups and Group Rings*, Tuscaloosa, AL, to appear.
- (9) Gaglione A.M. and Spellman, D., "Every 'universally free' group is tree-free," *Proc. Ohio State-Denison Conf. for H. Zassenhaus*, to appear.
- (10) Grätzer, G., *Universal Algebra*, Van Nostrand, 1968, USA.
- (11) Harrison, N., "Real length functions in groups," *Trans. Amer. Math. Soc.* 174(1972), 77 – 106.
- (12) Lyndon, R. C., "Length functions in groups," *Math. Scand.* 12(1963), 209– 234.
- (13) Morgan, J. W., " $\Lambda$ -trees and their applications," *Bull. Amer. Math. Soc.* 26(1992), 87 – 112.
- (14) Remeslennikov, V.N., " $\exists$ -Free groups and groups with a length function," Preprint.
- (15) Shelah, S., "Every two elementary equivalent models have isomorphic ultrapowers," *Israel J. Math.* 10(1971)224 – 233.

Department of Mathematics, U.S. Naval Academy, Annapolis, MD, 21402 – 5002  
*E-mail address:* amgma.usna.navy.mil

Philadelphia, PA19124 – 3036